

Trigonometric Functions \Leftrightarrow Hyperbolic Functions

Construction of relationships that transform hyperbolic functions into trigonometric functions.

The Pythagorean formula for a right triangle with hypotenuse “h” and side “a” adjacent to angle α and side “b” opposite angle α is:

$$h^2 = a^2 + b^2 \quad 33.20$$

For this triangle we have the trigonometric relations:

$$a = h \cdot \cos \alpha \quad b = h \cdot \sin \alpha \quad 33.21$$

Reshaping the Pythagorean formula gives:

$$h^2 = a^2 + b^2 \rightarrow b^2 = h^2 - a^2 = (h + a)(h - a) \rightarrow \left(\frac{h+a}{b}\right) \left(\frac{h-a}{b}\right) = e^\vartheta e^{-\vartheta} = 1 \quad 33.22$$

This is divided into the following hyperbolic functions:

$$e^\vartheta = \frac{h+a}{b} > zero \quad 33.23$$

$$e^{-\vartheta} = \frac{h-a}{b} > zero \quad 33.24$$

Where applying the trigonometric relations we obtain:

$$e^\vartheta = \frac{h+a}{b} = \frac{h+h \cdot \cos \alpha}{h \cdot \sin \alpha} = \frac{1+\cos \alpha}{\sin \alpha} \quad 33.25$$

$$e^{-\vartheta} = \frac{h-a}{b} = \frac{h-h \cdot \cos \alpha}{h \cdot \sin \alpha} = \frac{1-\cos \alpha}{\sin \alpha} \quad 33.26$$

From trigonometry we have:

$$\operatorname{tg} \left(\frac{\alpha}{2}\right) = \frac{1-\cos \alpha}{\sin \alpha} = \frac{\sin \alpha}{1+\cos \alpha} = \sqrt{\frac{1-\cos \alpha}{1+\cos \alpha}} \quad 33.27$$

Applying 27 we obtain the trigonometric angle α :

$$e^\vartheta = \frac{1+\cos \alpha}{\sin \alpha} = \frac{1}{\operatorname{tg} \left(\frac{\alpha}{2}\right)} = \frac{1}{\sqrt{\frac{1-\cos \alpha}{1+\cos \alpha}}} = \sqrt{\frac{1+\cos \alpha}{1-\cos \alpha}} \quad 33.28$$

$$e^{-\vartheta} = \frac{1-\cos \alpha}{\sin \alpha} = \operatorname{tg} \left(\frac{\alpha}{2}\right) = \sqrt{\frac{1-\cos \alpha}{1+\cos \alpha}} \quad 33.29$$

$$\alpha = 2 \operatorname{arctg} (e^{-\vartheta}) \quad 33.30$$

From these we obtain the fundamental formulas of the hyperbolic angle \emptyset :

$$\ln(e^\emptyset) = \ln \left[\frac{1}{\operatorname{tg}\left(\frac{\alpha}{2}\right)} \right] \rightarrow \emptyset = \ln \left[\frac{1}{\operatorname{tg}\left(\frac{\alpha}{2}\right)} \right] \quad 33.31$$

$$\ln(e^{-\emptyset}) = \ln \left[\operatorname{tg}\left(\frac{\alpha}{2}\right) \right] \rightarrow \emptyset = -\ln \left[\operatorname{tg}\left(\frac{\alpha}{2}\right) \right] \quad 33.32$$

In the unitary hyperbola $x^2 - y^2 = 1$ applying the functions $x = \operatorname{ch}\emptyset$ and $y = \operatorname{sh}\emptyset$ we get:

$$x^2 - y^2 = \operatorname{ch}^2\emptyset - \operatorname{sh}^2\emptyset = (\operatorname{ch}\emptyset + \operatorname{sh}\emptyset)(\operatorname{ch}\emptyset - \operatorname{sh}\emptyset) = e^\emptyset \cdot e^{-\emptyset} = 1 \quad 33.33$$

Breaking it down into two functions yields the hyperbolic cosine "ch \emptyset " and hyperbolic sine "sh \emptyset " functions:

$$\operatorname{ch}\emptyset + \operatorname{sh}\emptyset = e^\emptyset \rightarrow x = \operatorname{ch}\emptyset = \frac{e^\emptyset + e^{-\emptyset}}{2} \quad 33.34$$

$$\operatorname{ch}\emptyset - \operatorname{sh}\emptyset = e^{-\emptyset} \rightarrow y = \operatorname{sh}\emptyset = \frac{e^\emptyset - e^{-\emptyset}}{2} \quad 33.35$$

In 34 and 35 we have the fundamental properties of the hyperbolic functions.

Applying to the hyperbolic cosine ch \emptyset 34, the previous variables are obtained:

$$x = \operatorname{ch}\emptyset = \frac{e^\emptyset + e^{-\emptyset}}{2} = \frac{1}{2} \left(\frac{h+a}{b} + \frac{h-a}{b} \right) = \frac{h}{b} = \frac{h}{h \cdot \operatorname{sena}} = \frac{1}{\operatorname{sena}} = \operatorname{coseca} \quad 33.36$$

Applying to the hyperbolic sine sh \emptyset 35, the previous variables are obtained:

$$y = \operatorname{sh}\emptyset = \frac{e^\emptyset - e^{-\emptyset}}{2} = \frac{1}{2} \left(\frac{h+a}{b} - \frac{h-a}{b} \right) = \frac{a}{b} = \frac{h \cdot \operatorname{cosa}}{h \cdot \operatorname{sena}} = \frac{\operatorname{cosa}}{\operatorname{sena}} = \operatorname{cotga} \quad 33.37$$

Applying the hyperbolic cosine $x = \operatorname{ch}\emptyset = \operatorname{coseca}$ and the hyperbolic sine $y = \operatorname{sh}\emptyset = \operatorname{cotga}$ to the unitary hyperbola equation $x^2 - y^2 = 1$ we get:

$$x^2 - y^2 = \operatorname{ch}^2\emptyset - \operatorname{sh}^2\emptyset = \operatorname{cosec}^2\alpha - \operatorname{cotg}^2\alpha = 1 \quad 33.38$$

Which is a result of trigonometry.

With the relations of the hyperbolic cosine ch \emptyset and the hyperbolic sine sh \emptyset we can define the other relations between the trigonometric functions and the hyperbolic functions:

$$\operatorname{tgh}\emptyset = \frac{\operatorname{sh}\emptyset}{\operatorname{ch}\emptyset} = \frac{\frac{\operatorname{cosa}}{\operatorname{sena}}}{\frac{1}{\operatorname{sena}}} = \operatorname{cosa} \quad 33.39$$

$$\operatorname{cotgh}\emptyset = \frac{\operatorname{ch}\emptyset}{\operatorname{sh}\emptyset} = \frac{\frac{1}{\operatorname{sena}}}{\frac{\operatorname{cosa}}{\operatorname{sena}}} = \frac{1}{\operatorname{cosa}} = \operatorname{seca} \quad 33.40$$

$$\operatorname{sech}\emptyset = \frac{1}{\operatorname{ch}\emptyset} = \frac{1}{\frac{1}{\operatorname{sena}}} = \operatorname{sena} \quad 33.41$$

$$\operatorname{cosech}\emptyset = \frac{1}{\operatorname{sh}\emptyset} = \frac{1}{\frac{\operatorname{cosa}}{\operatorname{sena}}} = \frac{\operatorname{sena}}{\operatorname{cosa}} = \operatorname{tga} \quad 33.42$$

$$\operatorname{sech}^2\emptyset + \operatorname{tgh}^2\emptyset = \operatorname{sen}^2\alpha + \operatorname{cos}^2\alpha = 1 \quad 33.43$$

$$\operatorname{cotgh}^2\emptyset - \operatorname{cosech}^2\emptyset = \operatorname{sec}^2\alpha - \operatorname{tg}^2\alpha = 1 \quad 33.44$$

Construction of the already known relationships that transform the hyperbolic functions into the exponential form of a complex number.

Next, we will use Euler's formulas:

$$e^{i\alpha} = \cos\alpha + i\sin\alpha \qquad e^{-i\alpha} = \cos\alpha - i\sin\alpha \qquad 33.45$$

Reshaping the Pythagorean formula, we get:

$$h^2 = a^2 + b^2 = a^2 - (ib)^2 = (a + ib)(a - ib) \rightarrow \frac{(a+ib)}{h} \frac{(a-ib)}{h} = e^{\emptyset} e^{-\emptyset} = 1 \qquad 33.46$$

This breaks down into the following complex hyperbolic functions:

$$e^{\emptyset} = \frac{a+ib}{h} > zero \qquad 33.47$$

$$e^{-\emptyset} = \frac{a-ib}{h} > zero \qquad 33.48$$

For this triangle we have the trigonometric relations:

$$\frac{a}{h} = \cos\alpha \qquad \frac{b}{h} = \sin\alpha \qquad 33.49$$

Applying trigonometric relations, we get:

$$e^{\emptyset} = \frac{a+ib}{h} = \frac{a}{h} + i \frac{b}{h} = \cos\alpha + i\sin\alpha \qquad 33.50$$

$$e^{-\emptyset} = \frac{a-ib}{h} = \frac{a}{h} - i \frac{b}{h} = \cos\alpha - i\sin\alpha \qquad 33.51$$

To conform to Euler's formulas we must change the hyperbolic arguments to $\emptyset = i\alpha$ and thus we obtain the hyperbolic functions written as the exponential form of a complex number:

$$e^{\emptyset} = e^{i\alpha} = \cos\alpha + i\sin\alpha \qquad 33.52$$

$$e^{-\emptyset} = e^{-i\alpha} = \cos\alpha - i\sin\alpha \qquad 33.53$$

Calling the coseno $ch\alpha$ hyperbolic complex as:

$$x = ch\alpha = \frac{e^{i\alpha} + e^{-i\alpha}}{2} = \frac{1}{2}[(\cos\alpha + i\sin\alpha) + (\cos\alpha - i\sin\alpha)] = \cos\alpha \qquad 33.54$$

And naming the sine $sh\alpha$ hyperbolic complex as:

$$y = sh\alpha = \frac{e^{i\alpha} - e^{-i\alpha}}{2} = \frac{1}{2}[(\cos\alpha + i\sin\alpha) - (\cos\alpha - i\sin\alpha)] = i\sin\alpha \qquad 33.55$$

Applying the cosine $x = ch\alpha = \cos\alpha$ hyperbolic complex and the sine $y = sh\alpha = i\sin\alpha$ hyperbolic complex in the equation of the unit hyperbola $x^2 - y^2 = 1$ results:

$$x^2 - y^2 = ch^2\alpha - sh^2\alpha = \cos^2\alpha - i^2\sin^2\alpha = \cos^2\alpha + \sin^2\alpha = 1 \qquad 33.56$$

Which is a result of trigonometry.

With the relationships of the hyperbolic cosine $ch\alpha = \cos\alpha$ and the hyperbolic sine $sh\alpha = i\sin\alpha$ we can define the other relationships between complex trigonometric functions and complex hyperbolic functions.

Construction of relationships that transform hyperbolic functions into trigonometric functions similar to those that occur in Gudermannian functions.

The Pythagorean formula for a right triangle with hypotenuse “h” and side “a” adjacent to angle α and side “b” opposite angle α is:

$$h^2 = a^2 + b^2 \quad 33.57$$

For this triangle we have the trigonometric relations:

$$a = h \cdot \cos\alpha \quad b = h \cdot \sin\alpha \quad 33.58$$

Reshaping the Pythagorean formula gives:

$$h^2 = a^2 + b^2 \rightarrow a^2 = h^2 - b^2 = (h + b)(h - b) \rightarrow \left(\frac{h+b}{a}\right) \left(\frac{h-b}{a}\right) = e^\beta \cdot e^{-\beta} = 1 \quad 33.59$$

This is divided into the following hyperbolic functions:

$$e^\beta = \frac{h+b}{a} > \text{zero} \quad 33.60$$

$$e^{-\beta} = \frac{h-b}{a} > \text{zero} \quad 33.61$$

Where applying the trigonometric relations we obtain:

$$e^\beta = \frac{h+b}{a} = \frac{h+h \cdot \sin\alpha}{h \cdot \cos\alpha} = \frac{1+\sin\alpha}{\cos\alpha} \quad 33.62$$

$$e^{-\beta} = \frac{h-b}{a} = \frac{h-h \cdot \sin\alpha}{h \cdot \cos\alpha} = \frac{1-\sin\alpha}{\cos\alpha} \quad 33.63$$

From these we obtain the fundamental formulas of the hyperbolic angle β :

$$\ln(e^\beta) = \ln\left(\frac{1+\sin\alpha}{\cos\alpha}\right) \rightarrow \beta = \ln\left(\frac{1+\sin\alpha}{\cos\alpha}\right) \quad 33.64$$

$$\ln(e^{-\beta}) = \ln\left(\frac{1-\sin\alpha}{\cos\alpha}\right) \rightarrow \beta = -\ln\left(\frac{1-\sin\alpha}{\cos\alpha}\right) \quad 33.65$$

Denominating the hyperbolic cosine $ch\beta$ as:

$$x = ch\beta = \frac{e^\beta + e^{-\beta}}{2} = \frac{1}{2} \left(\frac{h+b}{a} + \frac{h-b}{a} \right) = \frac{h}{a} = \frac{h}{h \cdot \cos\alpha} = \frac{1}{\cos\alpha} = \sec\alpha \quad 33.66$$

And calling the hyperbolic sine $sh\beta$ as:

$$y = sh\beta = \frac{e^\beta - e^{-\beta}}{2} = \frac{1}{2} \left(\frac{h+b}{a} - \frac{h-b}{a} \right) = \frac{b}{a} = \frac{h \cdot \sin\alpha}{h \cdot \cos\alpha} = \frac{\sin\alpha}{\cos\alpha} = \tan\alpha \quad 33.67$$

Applying the hyperbolic cosine $x = ch\beta = \sec\alpha$ and the hyperbolic sine $y = sh\beta = \tan\alpha$ to the unitary hyperbola equation $x^2 - y^2 = 1$ we get:

$$x^2 - y^2 = ch^2\beta - sh^2\beta = \sec^2\alpha - \tan^2\alpha = 1 \quad 33.68$$

Which is a result of trigonometry.

With the relations of the hyperbolic cosine $ch\beta$ and the hyperbolic sine $sh\beta$ we can define the other relations between the trigonometric functions and the hyperbolic functions:

$$tgh\beta = \frac{sh\beta}{ch\beta} = \frac{\frac{sen\alpha}{\frac{1}{\cos\alpha}}}{\frac{1}{\cos\alpha}} = sen\alpha \quad 33.69$$

$$cotgh\beta = \frac{ch\beta}{sh\beta} = \frac{\frac{1}{\frac{\cos\alpha}{sen\alpha}}}{\frac{\cos\alpha}{sen\alpha}} = \frac{1}{sen\alpha} = cosec\alpha \quad 33.70$$

$$sech\beta = \frac{1}{ch\beta} = \frac{1}{\frac{1}{\cos\alpha}} = \cos\alpha \quad 33.71$$

$$cosech\beta = \frac{1}{sh\beta} = \frac{1}{\frac{\cos\alpha}{sen\alpha}} = \frac{sen\alpha}{\cos\alpha} = cotg\alpha \quad 33.72$$

$$sech^2\beta + tgh^2\beta = \cos^2\alpha + sen^2\alpha = 1 \quad 33.73$$

$$cotgh^2\beta - cosech^2\beta = cosec^2\alpha - cotg^2\alpha = 1 \quad 33.74$$